


PATHOPHYSIOLOGY OF BONES AND JOINTS


DENTAL MEDICINE
2024/2025

Eva Lovásová, MVD., PhD.
Department of Pathological Physiology
Faculty of Medicine, P. J. Šafárik University, Košice



1

DEFINITION OF BONE TISSUE



- Bone tissue forms most of the skeleton, the framework that supports and protects vital organs, bone marrow, acts as mineral reservoir for calcium, has function in acid-base homeostasis and, of course, allows movement.
- Bones are characterized their rigidity, hardness, and power of regeneration and repair.
- Strong but light weight, bone is a dynamic tissue. Throughout life, it is continually being broken down and formed.

2

FUNCTIONS OF BONE TISSUE

- Supports the muscles, teeth
- Protects the brain, spinal cord, heart, lungs and other inner organs
- Allows movement
- Maintains ion balance
- Helps with maintaining of acid-base balance
- Bone marrow produces blood cells




3

CLASSIFICATION OF BONES


ACCORDING TO POSITION

- **Axial skeleton**
 - Skull, vertebral column, ribs, sternum
- **Appendicular skeleton**
 - Skeleton of limbs



4

CLASSIFICATION OF BONES ACCORDING TO SHAPE



Long bones

- Longer than they are wide (e.g humerus)
- Consist of a long shaft with two bulky ends or extremities
- Primarily compact bone but may have a large amount of spongy bone at the ends or extremities

Short bones

- Cube shaped bones of wrist and ankle
- Consist mainly of spongy bone, which is covered by a thin layer of compact bone
- Bones that form within tendons(e.g patella)

Flat bones

- Thin, flattened and a bit curved (e.g sternum and most skull bones)

Irregular bones

- Bones with complicated shapes (e.g vertebrae,hip bone, maxilla, mandible)
- Primarily spongy bone that is covered with a thin layer of compact bone

5

CLASSIFICATION OF BONES ACCORDING TO MATURITY



Woven bone


- Immature bones
 - Fetal bones
 - Fracture healing
 - Areas surrounding tumors or inflammation
- Collagen fibers lay in disorganized manner

Lamellar bone

- Mature bone
- Regular parallel alignment of collagen fibers organized into sheets (lamellae)

6

CLASSIFICATION OF BONES ACCORDING TO STRUCTURE




- **Compact bone**
 - 85% skeleton
 - surface layer of short and flat bones and diaphysis of long bones
 - concentrically arranged bone lamellae - give the bone strength
 - between the lamellae in the cavities (lacunae) are bone cells osteocytes


- **Spongy bone**
 - 15% skeleton
 - a main part of the flat and short bones and epiphysis of the long bones
 - lamellae arranged in trabeculae, between which there are cavities, filled with red bone marrow

7

ANATOMY OF BONES



8



9

BONE CELLS

Osteoblasts

- Bone building cells that secrete bone matrix
- They lost the ability to divide by mitosis
- Secrete collagen and other organic components needed to build bone tissue
- The differentiation of osteoprogenitor cells into osteoblasts is accelerated by skeletal growth factors

Functions

- Role in formation of bone matrix
- Role in calcification (through the alkaline phosphatase enzymes)
- Synthesis of proteins

Osteoprogenitor cells

- Osteoblast precursors (preosteoblasts)
- Unspecialized cells derived from mesenchyme
- Undergo mitosis and develop into osteoblasts
- They also play a role in bone growth and healing, they can also differentiate into chondroblasts or fibroblasts.
- They are involved in the nutrition of bone tissue
- They are found in the periosteum, endosteum and in canals that contains blood vessels

10

BONE CELLS


Osteoblasts

- They synthesize the bone matrix - collagen
- Role in the deposition of inorganic substances in the bone matrix - calcification
- They synthesize growth factors - TGF β , GM-CSF, ...
- They produce enzymes - ALP

- They synthesize regulatory factors for remodeling
 - **RANKL (RANK ligand)** → binds to the RANK receptor (receptor activator of nuclear factor kappa B), which is found on osteoclast precursors → osteoclast precursors change into mature osteoclasts → **ACTIVATION OF BONE RESORPTION**
 - **Osteoprotegerin (OPG)** - prevents the binding of RANKL to the RANK receptor on osteoclasts → **INHIBITION OF BONE RESORPTION** (it is also the RANK receptor)


11

RANKL/RANK/OSTEOPROTEGERIN



12

BONE CELLS



Osteocytes

- Bone maintenance
- Small star-shaped cells placed in lacunae communicating with each other through protrusions
- Derived from mature osteoblasts
- Produced by RANKL and OPG
- Produce also **sclerostin**, protein that has anti-anabolic effects on bone formation – inhibits function of osteoblasts
- Important for the existence of the extracellular matrix, they have a low synthetic capacity, they also participate in resorption
- maintain the bone matrix, regulate bone metabolism, participate in the transport of substances, especially ions (calcium, phosphates) between the bone and blood plasma
- their protrusions serve as mechanoreceptors, transmit mechanical stimulation from the bone surface to osteocytes, which on the basis of this information in cooperation with osteoblasts and osteoclasts activate bone formation or resorption

13

BONE CELLS




Osteoclasts

- Large multinucleated cells
- They are involved in bone resorption
- Derived from the same hematopoietic lineage as monocytes and macrophages
- They contain a large number of lysosomes filled with hydrolytic enzymes
- They bind to the bone surface via receptors for adhesive molecules. At the point of contact, the membrane osteoclasts form a typical wavy edge by which they adhere to the bone surface. This increases the size of the contact cells and determines the area of bone resorption and protects the surrounding tissue.
- Formation of **hydrochloric acids (HCl)**, which acidifies the area of resorption, and breaks down the mineral matrix
- They release the protease **cathepsin K** from lysosomes, which breaks down bone matrix proteins, especially collagen I.
- Ions formed by the decomposition of hydroxyapatite (calcium, phosphate) and collagen fragments are absorbed by osteoclasts (endocytosis), transported in vacuoles to the opposite pole of the cell (transcytosis) and excreted into the extracellular fluid.

14

BONE MATRIX

- **Inorganic components (cca 70 %)**
 - Hydroxyapatite $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ (99 %)
 - Other - CaHPO_4 , MgHPO_4 , CaCO_3
- **Organic components (cca 25 %)**
 - Collagen 1 (90 – 95 %)
 - Osteocalcin (protein)
 - Osteonectin
 - Proteoglycans
 - Glykoproteins
 - Sialoproteins
 - Lipids
- **Water (cca 5 %)**



15

OSTEOGENESIS (BONE GROWTH)

Ossification (osteogenesis) - process of formation of new bone by osteoblasts.

Types of osteogenesis

1. Intramembranous ossification

- Laying down of bone into the primitive connective tissue (mesenchyme) resulting in the formation of bones (skull, clavicle, mandible).
- Healing process of fractures

2. Endochondral ossification

- Cartilage acts as a precursor (e.g., femur, tibia, humerus, radius)
- Growing of the length of long bones
- Fracture healing
 - *Perichondrial ossification* - on the surface of the cartilage model
 - *Enchondrial ossification* - inside the cartilaginous model

Steps of osteogenesis


1. Synthesis of extracellular organic matrix (osteoid)
2. Matrix mineralization leading to the formation of bone
3. Remodeling of bone by the process of resorption and reformation

16

ENDOCHONDRAL OSSIFICATION

Steps

1. Development of cartilage model
2. Growth of cartilage model
3. Development of the primary ossification center
 - fetal development of bones
 - a few short bones begin their primary ossification after birth
4. Development of the secondary ossification center
 - after birth
 - forms the epiphyses of long bones and the extremities of irregular and flat bones
5. Formation of articular cartilage and epiphyseal plate
 - after reaching of skeletal maturity (14–18 years of age), all of the cartilage is replaced by bone, fusing the diaphysis and both epiphyses together (epiphyseal closure).



17

BONE REMODELING

- a lifelong process - old bone is removed from the skeleton (bone resorption), and new bone is added (bone formation)

Function

- maintaining of normal function, structure and mineral homeostasis of bone
- healing of injuries like fractures but also microdamage which occurs during normal activity

The average lifespan of each remodeled unit in humans is 2–8 months.

In the young skeleton, the amount of resorbed bone is proportional to the newly formed - balanced process.

Up to the third decade – positive balance.

In the third decade - bone mass is at its maximum, and this is maintained with small variations until the age of 50.

After the age 50 resorption predominates and the bone mass begins to decrease.

Bone remodeling increases in perimenopausal and postmenopausal women.

18

REMODELING UNIT

Osteoclasts

- resorbing of bone
- derived from mononuclear precursor cells

Bone resorption depends on osteoclast secretion of hydrogen ions and cathepsin K enzyme. H^+ ions acidify the resorption compartment to dissolve the mineral component of bone matrix. Cathepsin K digests the proteinaceous matrix, which is mostly composed of type I collagen.

Osteoblasts

- bone formation
- stimulated by growth hormone, thyroid hormones, estrogens, androgens

19

REMODELING UNIT

RANK

The cell surface receptor RANK (receptor activator of NF κ B) activate osteoclast precursor cells to develop into fully differentiated osteoclasts when RANK is activated by its RANK ligand (RANKL). RANKL is produced mainly by marrow stromal cells and osteoblasts.

Osteoprotegerin

Osteoprotegerin (OPG), also known as osteoclast inhibiting factor (OCIF) or osteoclast binding factor (OCIF), is a key factor inhibiting the differentiation and activation of osteoclasts, Osteoprotegerin inhibits the binding of RANK to RANKL and inhibits the activation of osteoclasts.

Abnormalities in the balance of RANKL/RANK/OPG system lead to the increased bone resorption that underlies the bone damage of postmenopausal osteoporosis, Paget's disease, bone loss in metastatic cancers, and rheumatoid arthritis.

20

REMODELING PHASES

1. Quiescent Phase

bone is at rest

2. Activation Phase

activation of the bone surface to resorption

activation of osteoclast precursors - differentiation, migration, and fusion of the large multinucleated osteoclasts. These cells attach to the mineralized bone surface and initiate resorption by the secretion hydrogen ions and cathepsin K, which degrade bone matrix.

3. Resorption Phase

the osteoclasts dissolve the mineral matrix

4. Reversal Phase


bone resorption transitions to bone formation

5. Formation Phase

osteoclasts have resorbed a cavity of bone, they detach from the bone surface and are replaced by the osteoblast lineage which in turn initiate bone formation

6. Mineralization Phase

21



22

REGULATION OF SKELETAL METABOLISM


Parathyroid Hormone

- Peptide hormone (84 amino acids), produced by parathyroid glands
- regulates serum calcium and phosphorus concentrations through its receptor-mediated, combined actions on bone, intestine, and kidney
- high levels of PTH increase osteoclastic bone resorption
- low levels increase osteoblastic bone formation
- PTH receptor mainly at the osteoblasts

Calcitonin

- Peptide hormone (32 amino acids) produced by parafollicular C-cells of thyroid gland
- inhibit osteoclast-mediated bone resorption

23



24

REGULATION OF SKELETAL METABOLISM

Vitamin D (calcitriol)

Intestinum

- increases intestinal calcium absorption by increasing calcium uptake (calbindin)

Bone

- promotes the mineralization of osteoid and causes bone resorption by mature osteoclasts

- regulates the expression of several bone proteins (osteocalcin)


Kidney

- decreases calcium and phosphorus excretion

Blood

- increases calcium and phosphorus levels

25



26

REGULATION OF SKELETAL METABOLISM

Other hormones

Parathyroid hormone-related protein (PTHrP)

- Function as parathyroid hormone, produced by cancers but also physiologically E.g. tooth eruption - local bone resorption

Androgens

- Anabolic effect through the stimulation of the osteoblast receptors
- In childhood activate growth factors – increase bone density
- On the end of puberty activate epiphyseal closure

Estrogens

- After puberty - epiphyseal closure, but stimulate bone remodeling – after menopause deficit causes osteoporosis

Glucocorticoids

- Bone cell differentiation during development

Insulin

- Stimulates matrix synthesis together with IGF-1

Growth hormone

- Stimulates osteoblasts activity

Thyroid hormones

- Stimulate growth resorption but also formation, stimulate the synthesis of osteoid matrix and its mineralization

Growth factors and cytokins (IGF-1)

27

REGULATION OF SKELETAL METABOLISM

Decrease bone resorption

- Calcitonin
- Estrogens

Increase bone resorption

- PTH/PTHrP
- Glucocorticoids
- Thyroid hormones
- High-dose vitamin D

Increase bone formation

- Growth hormone
- Vitamin D metabolites
- Androgens
- Insulin
- Low-dose PTH/PTHrP
- Progestogens

Decrease bone formation

- Glucocorticoids

28

REGULATION OF SKELETAL METABOLISM

Other factors

Growth factors and cytokines

- Insulin-like growth factors I and II (IGF-I and II) - Increase osteoblast number and activity, increase collagen synthesis
- Interleukin 1 (IL-1) - Increases bone resorption

Other factors

- Genetic predisposition - The amount of bone tissue is partially inherited. The difference is between the races, black people have a stronger skeleton than whites, the lowest bone tissue amount have Asians.
- Movement - Lack of exercise increases bone breakdown, while regular movement promotes bone formation
- Nutritional factors - Malnutrition leads to bone loss. Calcium in the diet is essential for bone mineralization. Smoking, coffee, alcohol, lots of salt in the diet increase the risk of osteopenia

29

ANATOMY AND PHYSIOLOGY OF JOINTS

Joint (articulation) - the place of connection of two or more bones.

The role of the joints is to allow movement and support the stability of the skeleton.




30

CLASSIFICATION OF JOINTS

ACCORDING TO MOBILITY

- immovable (*synarthrosis*)
- slightly movable (*amphiarthrosis*)
- Freely movable (*joints, diarthrosis*)




31

CLASSIFICATION OF JOINTS

Movable (synovial) joints

- They are formed by articulating bones
- The surfaces of the bones that are in contact are covered with hyaline cartilage, which exactly copies the shape of the joint. Its thickness is 0.5 to 6 mm.
- Between the joint surfaces - cavity filled with synovial fluid. The cavity can be divided into two parts using the so-called articular disc (meniscus).
- The whole joint is surrounded by a fibrous capsule, the inner layer - synovial membrane produces synovial fluid and the outer layer, the fibrous membrane, strengthens the joint.



32

CLASSIFICATION OF JOINTS

Immovable joints

Fibrous



- Syndesmosis - bones held together by an interosseous membrane. The middle radioulnar joint and middle tibiofibular joint are examples of a syndesmosis joint
- Sutura - is a junction of flat bones, e.g. skull bones in children
- Gomphosis - where the teeth articulate with their sockets in the maxilla (upper teeth) or the mandible (lower teeth).

Cartilaginous


- Synchondrosis - the bones are connected by hyaline cartilage - the joint between the diaphysis and epiphysis of a growing long bone
- Symphysis - pubic symphysis

Bone

- Synostosis - fusion of two or more bones - e.g. the connection of the five sacral vertebrae forming the sacrum



33



PATHOPHYSIOLOGY OF BONES

34

OSTEOPOROSIS


- common metabolic disorder of the skeleton
- bone mineral density is reduced
- the bone microarchitecture is disrupted (perforation of trabecular plates)
- the amount and variety of noncollagenous proteins in bone is altered

Leads to

- increased risk of fracture

Causes

- **Primary** – postmenopausal, idiopathic juvenile, senile
- **Secondary** – nutrition (anorexia), endocrine (Cushings, hyperprolacinemia), drug (corticoids), malignancy, chronic diseases (diabetes mellitus), genetic (Klinefelter sy., Turner sy.), idiopathic



35

OSTEOPOROSIS

Risk factors


- Age
- Female mainly postmenopausal
- Family history
- Sedentary lifestyle
- Defficiency of calcium or vitamin D
- High protein, alcohol, caffeine intake
- Smoking
- Hyperparathyroidism
- Other diseases (diabetes mellitus, celiac disease)
- Drugs (Al-containing antacids)

Mechanisms

- \downarrow estrogens \rightarrow \uparrow cytokines \rightarrow \uparrow RANKL \rightarrow \uparrow activity of osteoclasts
- \uparrow sclerostin \rightarrow \downarrow activity of osteoblasts



36



37

OSTEOPOROSIS

Clinical signs

- „silent killer of bones“ – long time without signs, pain
- decrease of height
- fractures of bones – hip, humerus,... - bones become brittle and there is a risk of fracture even when exposed to a disproportionately low intensity of mechanical energy, e.g. fracture of the femoral neck when falling on its side, spinal fractures when lifting heavy objects, wrist fractures when banging on a table, etc.
- deformities in the thoracic spine with back pain
- reduction in body height
- wedging and collapse of vertebrae
- kyphosis
- hump

38

OSTEOMALACIA

- inadequate mineralization of bones

Causes

- deficiency of vit. D – ↓ absorption of calcium from the intestine
- phosphate deficiency due to renal diseases or ↓ of intestine absorption

Clinical signs

- In the beginning - non-specific symptoms: muscle weakness, fatigue, muscle, bone and joint pain
- Diffuse joint and bone pain - lumbosacral spine, pelvis, ribs, forelegs and plexus areas of the limbs. Pain most often occurs with movement
- Bones are sensitive to touch.
- Later - deformities, mainly in the area of the chest and lower limbs
- Fractures - less common than in osteoporosis
- In the advanced stage of the disease - immobility, pain and hypersensitivity to touch



39

RICKETS


- inadequate mineralization of bones in children whose epiphyses have not yet fused

Causes


- deficiency of vit. D
- Vit. D resistant rickets
 - phosphate deficiency
 - mutation of vit. D receptor

Clinical signs

- deformation, widening of the bones
 - In newborns – deformation of skull
 - In older children - the lower limbs
- As a result of insufficient mineralization, the growth plates are widening, which is manifested by swelling most often on the ankles, wrists and sternal ends of the ribs
- Growth is slowed, muscles are hypotonic, gait is slow "duck"



40



41

PAGET'S DISEASE

- Osteitis deformans
- disorder of bone remodeling
- the osteoclasts become abnormally activated, and produce a bizarre and irregular pattern of resorption, to which there is usually an intense osteoblastic response with irregular new bone formation often in the form of woven bone
- increased bone density, but because of the irregular architecture, bone strength is decreased and pathologic fractures may occur.

Causes

- hereditary?
 - genetically determined increased sensitivity of osteoclasts to activation by RANKL
- environmental factors – virus infection
 - viruses from the paramyxoviridae family

Clinical signs

- Enlargement and deformation of bones – face, long bones
- Pain, fractures that bleed heavily
- CVD disorders
- Tooth loss
- Hearing disorders, vestibular disorders, dementia
- Bone sarcomas

42



PATHOPHYSIOLOGY OF JOINTS

43

OSTEOARTHRITIS


- Degenerative changes of joints

Causes

- Unknown
- Genetic predisposition
- Environmental factors
- Obesity
- Mechanical stress

Classification

- **Primary arthrosis**
 - causes are unknown
 - genetic predisposition (quality of cartilage, strength of ligaments)
 - risk factors - obesity, uneven loading of joints (incorrect footwear), long-term mechanical stress (sports, dancing).
- **Secondary arthrosis**
 - the consequence of another primary joint disorders, e.g. in some metabolic diseases (alkaptonuria, hemochromatosis), in some congenital disorders (dysplasia of the hip joint, hypermobility of the joints), after injuries (fractures, bleeding in the joint), in inflammatory diseases (rheumatoid arthritis).



44



OSTEOARTHRITIS

Pathogenesis

Slow progressive degeneration of articular cartilage, cartilage is damaged, loses elasticity, thins, joint space narrows, bones come into contact, causing pain and limitation of movement. Outgrowths (osteophytes) form on the edges of the joints. Synovial fluid production increases. Although arthrosis is not primarily an inflammatory disease, due to changes and increased irritation in the joint, local inflammation occurs secondary to it, which contributes to pain and limitation of movement.

Clinical signs

- Pain
- Difficulty to initiate movement - morning stiffness
- Joint crepitation



45

RHEUMATOID ARTHRITIS

- Chronic autoimmune systemic disease
- Synovial inflammation and destruction of the bone architecture
- Women vs. Men 3 : 1

Causes

- Autoimmune
 - Seropositive form - rheumatoid factor (antibody)
 - Seronegative form
- T-cells, B-cells, antibodies, macrophages, cytokines (TNF α , IL-1),
- Genetic predisposition - HLA-DR4-Dw4 alleles
- Production of cytokines - activate osteoclasts
- Trigger factor - infection (virus, bacteria, fungi)
- Chronic parodontitis is risk factor





46


RHEUMATOID ARTHRITIS

Clinical signs

- Joints
 - Attacks of symmetric joint pain and morning stiffness
 - Inflammation and later fibrosis cause movement limitation
 - Later deformation, dislocation of joints
- Extra-articular sign
 - Fatigue, weakness, anorexia, weight loss
 - Rheumatoid nodules
 - Fibrosis of lungs
 - Atherosclerosis



47



48

	Osteoarthritis	Rheumatoid Arthritis
Site(s) affected	Localized to joint	Articular, systemic and extra-articular manifestation
Pathogenesis	Biomechanical, leads to loss of cartilage matrix	Autoimmune response leads to joint destruction
Symptoms	Pain Stiffness <20 minutes Limited motion	Pain Joint swelling Stiffness >1 hour Limited motion
Inflammation	Usually limited, may be present in advanced disease	Chronic
Osteophytes	Usually present	Absent
Rheumatoid factors	Absent	Frequently present

OSTEOARTHRITIS
DEGENERATIVE DISEASE
MORNING STIFFNESS LASTING LESS THAN 30 MINUTES
HEBERDEN'S NODES
CARTILAGE LOSS
ASYMMETRICAL

RHEUMATOID ARTHRITIS
AUTOIMMUNE DISEASE
MORNING STIFFNESS LASTING MORE THAN 30 MINUTES
EXTRA-ARTICULAR INVOLVEMENT
INFLAMED SYNOVIUM
SYMMETRICAL

WWW.MEDCOMIC.COM © 2016 JORGE MUNIZ

49

ANKYLOSING SPONDYLITIS


- Systemic inflammatory disease of the joints characterized by stiffness and fusion (ankylosis) of the joints of the spine
- Especially in young people, men > women

Causes

- Autoimmune disease ?? with a genetic predisposition (HLA-B27).

Pathogenesis

- Inflammation affects the ligament-cartilage, cartilage and bone parts of vertebrae → erosions, especially in the cartilage, are compensated by proliferation of fibroblasts, which synthesize collagen, whose fibers fill erosions, which subsequently calcify and ossify → joints fuse and lose mobility.





50

ANKYLOSING SPONDYLITIS

Clinical signs

- Initially pain in the lumbar-sacral region, stiffness and pain when breathing, pain in the lower back is typical in the morning, or after a long rest, the movement is relieved
- Later, the patient has trouble sitting upright and turn. The spine changes shape, the typical lower curvature of the spine disappears and the upper part of the spine is tilted forward → breathing problems
- Sometimes the joints of the limbs (more often in women)
- Extra-articular manifestations - lung fibrosis, cardiomegaly, amyloidosis and others.



51

GOUT (ARTHRITIS URICA)

Cause - ???

- congenital - ↑ production of uric acid - enzyme defects
- ↓ excretion of uric acid - defect of renal transporters
- acquired - ↑ production of uric acid - obesity, alcohol
- high intake of food containing purins, leukaemia, haemolysis, cytostatic drugs, Gierke's disease
- ↓ excretion of uric acid - kidney disease
- drugs



Clinical symptoms

- hyperuricemia
- arthritis - acute arthritis
- chronic arthritis - tophi
- chronic uric acid interstitial nephropathy
- uric acid nephrolithiasis (kidney stones)



52


PATHOPHYSIOLOGY OF THE TEMPOROMANDIBULAR JOINT

- Jaw (temporomandibular) joint
- Opening and closing of mouth, chewing, yawning, talking ...
- It is one of the complex joints
- During the day, it is opened about 1800 times - it is one of the most used joints in the body



Causes of disorders

- dislocation (luxatio) - injury, yawning, eating of big pieces ...
- fracture - injury
- developmental defects
- swelling - inflammation, after anesthesia
- tumor
- osteoarthritis
- Convulsions - epilepsy, tetanus, diseases of the salivary glands ...
- Gritting of teeth (bruxism) during stress
- ...




53

PATHOPHYSIOLOGY OF THE TEMPOROMANDIBULAR JOINT

Clinical signs

- Joint pain, toothache
- Migraine headaches
- Cervical spine pain
- Chewing and swallowing disorders
- Tension, cramps
- With osteoarthritis - morning stiffness, creaking in the joint
- Tingling in the ears (tinnitus), vestibular symptoms
- "Ear" pain
- „Eye“ pain
- Stress, depression, anxiety



54

THANK YOU FOR YOUR ATTENTION



55